

УДК 616.711-007.43-089.8-072.1-71(045)

The current state of endoscopic disc surgery: review of controlled studies comparing full-endoscopic procedures for disc herniations to standard procedures¹

Christof Birkenmaier¹, Martin Komp², Hansjörg F. Leu³, Bernd Wegener¹, Sebastian Ruetten²

¹ Department of Orthopaedics, University of Munich (LMU), Grosshadern Campus, Munich. Germany

² Department of Spine Surgery and Pain Therapy, Center for Orthopaedics and Traumatology, St. Anna-Hospital Herne, University of Witten/Herdecke, Herne. Germany

³ The Bethania Spine Base, Bethania Hospital, Zurich. Switzerland

Background: Endoscopic spinal surgery is increasingly popular because it minimizes access trauma and hastens recovery from the intervention. Objective: To assess the clinical outcomes and complication rates of full-endoscopic disc surgery compared to the microsurgical standard procedures. Methods: A PubMed and Embase search was performed, considering entries up to January 2013. Only 5 controlled trials of 504 articles could finally be considered for evaluation. Results: Overall, the endoscopic techniques had shorter operating times, less blood loss, less operative site pain, and faster postoperative rehabilitation / shorter hospital stay / faster return to work than the microsurgical techniques. All 5 studies had fewer complications with the endoscopic technique. Conclusions: The studies show that full-endoscopic disc surgery can achieve the same clinical results in symptomatic cervical and lumbar disc herniations as the microsurgical standard techniques. Key words: neuropathic pain, disc herniation, cervical, lumbar, endoscopic, endoscopy, review.

Предпосылки: популярность эндоскопической хирургии позвоночника возрастает, так как она минимизирует травматичность доступа и ускоряет выздоровление после вмешательства. Цель: оценить клинические результаты и частоту осложнений после эндоскопических операций на межпозвоноковых дисках по сравнению с микрохирургическими стандартными операциями. Методы: был проведен информационный поиск с помощью электронно-поисковых систем PubMed и Embase с учетом поступлений до января 2013 г. Только 5 контролируемых исследований из 504 статей были в конечном итоге отобраны для изучения. Результаты: в целом, эндоскопические операции характеризовались более коротким временем вмешательства, меньшей кровопотерей, меньшей интенсивностью боли в области операционной раны и более быстрым восстановлением / менее продолжительным пребыванием в больнице / более быстрым возвращением к труду, чем после микрохирургических операций. Во всех пяти работах было отмечено меньшее количество осложнений в результате эндоскопических операций. Заключение: результаты исследования показывают, что полностью эндоскопические операции на межпозвоноковых дисках могут обеспечить такие же клинические результаты при симптоматических грыжах в шейном и поясничном отделах позвоночника, как и микрохирургические стандартные операции. Ключевые слова: нейропатическая боль, грыжа межпозвонокового диска, шейный и поясничный отдел, эндоскопия, обзор.

Key words: neuropathic pain, disc herniation, cervical, lumbar, endoscopic, endoscopy, review

¹ This article was published in the Pain Physician Journal 2013; 16; 335–344

Introduction

Neuropathic pain caused by cervical or lumbar disc herniations is among the most common reasons for which patients seek specialist treatment. While guidelines in many countries stress the point that surgery should be reserved for cases with fresh motor deficits or cauda equina syndrome, **the reality is that most disc surgeries, regardless of the technique used, are performed for nerve root pain (sciatica or cervicobrachialgia).** It needs to be **recognized that minimally invasive procedures with the goal to remove the herniation causing the problem are the logical next step after conservative measures and image-guided injection techniques have failed to provide adequate pain relief.**

Endoscopic surgery attempts to bridge the gap between injection techniques and open surgery in as much as it attempts to perform the decompression required via the most minimized surgical approach possible, which is the placement of an instrument of just a few millimeters in diameter over the spinal needle that otherwise would have been used to perform a selective nerve root block or a different type of injection. Early on, there already had been indications for better outcomes with less invasivity [1]. Endoscopic disc surgery was pioneered in the late 1980s and in the early 1990s, but for a number of reasons did not break into the mainstream of spinal therapies at that time [2–7].

Recent years have again seen growing interest in spinal endoscopy as well as the development of new anatomical approaches. The technically demanding field of spinal pain treatments is late in adopting this technology and this was made possible only by technical advances in the field of cameras, coaxial working sleeves, optics, video processing equipment, radiofrequency devices, and others. After all, joint arthroplasty had been firmly established for many years before spinal arthroplasty ever became a viable treatment option. Similarly, endoscopic techniques have become the gold standard for a large number of conditions in orthopedics, gynecology, anesthesiology, and surgery, while in spinal surgery they are still considered outsider procedures by many.

It therefore appears to be a suitable point in time to review the available studies on endoscopic disc surgery and to compare spinal endoscopy to the respective standard surgical procedures with regards to outcome and complications. This review is not a systematic review for a very practical reason. The scientific evidence for the superiority of microdiscectomy over conservative therapy or over standard open discectomy is still very weak, even though microdiscectomy represents the currently accepted gold standard as far as surgical treatments are concerned [8–11]. A review of trials

comparing newer procedures to microdiscectomy on the background of their relative levels of evidence would therefore inevitably come to the conclusion that there still is insufficient evidence to allow for any definitive conclusions.

The goal of this review is to investigate whether controlled studies exist that allow for the objective comparison of full-endoscopic spinal procedures to the respective gold standard procedures with regards to outcome and complications as the paramount clinical parameters on which treatment decisions are to be based. The scope of this review is limited to endoscopic disc surgery, primarily because the surgical treatment of symptomatic disc herniations is a very frequently performed spinal procedure. It also represents by far the most common spinal condition treated by means of endoscopy **and for which established standard procedures, such as microdiscectomy or keyhole foraminotomy exist, against which endoscopic procedures can be compared.** Papers primarily focusing on laser disc decompression (without targeted disc fragment extraction), which is **sometimes performed under endoscopic visualization,** were to be excluded from the search strategy. Different from Nellensteijn et al. [12] systematic review on transforaminal endoscopic disc surgery, studies on simple endoscopic decompression of the intradiscal space and/or indirect endoscopic decompression of the spinal canal by means of the «in-out-technique» were to be excluded. These techniques no longer represent the current standard of endoscopic disc surgery, which is **the direct extraction of disc fragments from the epidural space/the foramina and the direct decompression of neural structures under full visual control.**

Another fundamental difference from the recent review by Nellensteijn et al. [12] is that our review is not limited to the transforaminal approach. While transforaminal was the first endoscopic approach for accessing the disc space and the ventral epidural space, interlaminar approaches have been established for the lumbar **and for the cervical spine.** A far lateral transforaminal approach has been described for the lumbar spine and an anterior transdiscal approach exists for the cervical spine. For the purpose of this review, we therefore must also define what we consider «truly endoscopic» in the context of spinal applications, since there is a longstanding confusion of termini technici with regards to spinal endoscopy.

In the majority of other surgical specialties, endoscopy implies the use of a thin tubular optical and surgical device that is passed completely percutaneously by means of a stab incision. This is very different from using tissue dilators of increasing diameter in

order to introduce tubular mini-retractor systems for creating a small, but nevertheless open access portal, sometimes **even in combination with an operating microscope**. Some experimental studies suggest that the use of tubular retractor systems is less traumatic than microdiscectomy **on the basis of intraoperative electromyogram measurements and postoperative serum cytokine levels** [13, 14]. However, recent randomized controlled trials (RCTs) indicate that such tubular mini-retractor systems seem to offer no clinically relevant advantage over standard microdiscectomy and that they may have the potential of higher complication rates compared to either the microsurgical or the older open technique [15, 16].

The authors of this review distinguish the following 3 technical approaches to disc surgery as being separate entities.

Microdiscectomy

The use of a Caspar retractor or similar device and of an operating microscope to perform disc surgery through a small skin incision of only a few centimeters. In the lumbar spine, this is currently considered the gold standard.

Tubular Discectomy

The use of tissue dilators and of a tube system through a minimized incision of less than 2 cm together with an operating microscope. The most common example would be the MetriX tube system. The term «micro-endoscopic discectomy» is frequently used as a synonym. Constant irrigation is not generally used. Overall, this technique has much in common with the microsurgical approach but it reduces the access trauma by means of a blind transmuscular dilatation as compared to **an open, visually controlled muscle dissection** from the spinous process/lamina. The Destandeu endoscopic system is a special variation that also falls into this category. What differentiates this technique from the MetriX tube system is mainly the use of an endoscope/monitor system and a blunt, single-step dilatation of the perispinal muscles.

Endoscopic Disc Surgery

This entails the use of a thin tubular device that contains the optical system and a working channel. It is introduced completely percutaneously through a stab incision. Usually, a spinal needle–guide wire technique is used to secure the controlled trajectory of a blunt trocar to the desired spinal region. The working sleeve is then passed over the trocar after removal of the guide wire. Visualization is always achieved by means of a connected video camera and monitor system. The terms «percutaneous endoscopic discectomy» or «fullendoscopic discectomy» **have been used synonymously**. A monoportal technique is standard and

surgery is performed under constant saline irrigation. For the purpose of this review, it was decided to focus exclusively on category 3, truly endoscopic disc surgery, also known as «full-endoscopic disc surgery». These 3 surgical techniques are to be distinguished from pure epiduroscopic adhesiolysis, which has its own merits, but does not remove herniated disc material or other physical sources of direct nerve root compression [17–25].

Methods

PubMed and Embase database searches were performed using the following search strategy:

(endoscopic OR endoscopy) AND (disc OR discal OR disk OR diskal) AND (cervical OR lumbar OR lumbosacral) NOT laser. Database entries up to January 31, 2013, were considered. The returned results were screened and assigned to one of the following groups:

1. RCT
2. controlled studies (CS)
3. comparative studies
4. case series
5. case reports
6. review articles
7. technical articles, anatomical studies, reports on personal experience and letters
8. articles on laparoscopic spinal fusion
9. unrelated publications.

Only articles categorized as RCTs and CSs were considered for this review.

Results

Using the above search strategy, 504 references were retrieved. **Twelve of these references were classified as RCTs** [10, 13, 15, 16, 26–32] and another 15 were classified as CSs [33–47]. **The majority of references were categorized as case series** (169), studies on tubular discectomy (113), articles on laparoscopic fusion surgery (38), review articles (79), or otherwise unrelated to the topic (95).

Of the 12 RCTs, only 5 were using a true endoscopic technique as defined above and only 4 of these 5 trials had a standard procedure as a control group [29–32]. Of the 16 CSs, only 6 [36, 37, 39, 44, 47] were using a true endoscopic technique as defined above. Two of these latter 6 studies used an old intradiscal technique [4, 39], **one compared endoscopic biopsy for spondylodiscitis to CT-guided biopsy** [44] and **2 did not employ a control group as a standard technique** [36, 47]. The remaining controlled study had a retrospective, non-randomized study design and used intradiscal decompression in addition to direct visually controlled fragmentectomy [37], allowing for discussion of its

Table 1

Study Characteristics

Study	type	pro/ retro	herniation type	n total	m : f	Randomiz. mode	endoscopic procedure	n	comparison procedure	n	FU	FU Rate
Ruetten et al, 2008 [20]	RCT	pro	cervical lateral	200	68 : 132	not blinded, alternating assignment	endoscopic posterior foraminotomy	100	microsurgical ACDF (PEEK cage, no plate)	100	2y	88 %
Ruetten et al, 2008 [21]	RCT	pro	lumbar (median, lateral, extraforaminal)	200	84 : 116	not blinded, alternating assignment	endoscopic interlaminar or transforaminal sequestrectomy	100	microsurgical sequestrectomy (paramedian or lateral)	100	2y	89 %
Ruetten et al, 2009 [22]	RCT	pro	lumbar recurrent	100	56 : 44	not blinded, alternating assignment	endoscopic interlaminar or transforaminal sequestrectomy	50	microsurgical sequestrectomy (paramedian approach)	50	2y	87 %
Ruetten et al, 2009 [23]	RCT	pro	cervical	120	43 : 77	balanced block randomization, not blinded	anterior transdiscal endoscopic decompression	60	microsurgical ACDF (PEEK cage, no plate)	60	2y	86 %
Lee et al, 2009 [28]	CS	retro	lumbar recurrent	54	38 : 16	not randomized	endoscopic transforaminal sequestrectomy and disc decompression	25	microsurgical sequestrectomy (paramedian approach)	29	34 m	n. a.

Note. Type specifies randomized controlled trial (RCT) vs. controlled study (CS), pro/retro specifies prospective vs. retrospective study design, n total = total number of patients studied in both groups, m : f = male vs. female ratio, FU = time of follow up specified in years (y) or months (m), FU Rate specifies which percentage of patients were available at final follow up, ACDF = anterior cervical decompression and fusion, PEEK = poly-ether-ether-ketone, n. a. = not applicable.

results only with certain limitations. In summary, our search retrieved 4 RCTs that each compare a modern full-endoscopic technique for the treatment of cervical or lumbar disc herniations to an established standard microsurgical procedure. One additional CS can be considered only with clear limitations because of its retrospective, non-randomized study design and its particular surgical technique.

Established and standardized parameters for assessing clinical and radiological outcomes were used in all of these trials; the numbers of patients included are high enough to allow for clinically relevant conclusions. All 5 studies originate from very experienced groups of investigators, with all 4 RCTs having the same authors. The characteristics and the relevant results of these trials are displayed in Tables 1 and 2. As can be seen from Table 1, the studies that were retrieved differ with regards to spinal region (cervical, lumbar), type of approach (transforaminal lumbar, interlaminar lumbar, interlaminar cervical, anterior transdiscal cervical), and type of herniation (recurrent, primary), as well as with regards to the respective comparison procedure. This rendered it unreasonable to use the source data of the individual trials as the basis for a meta-analysis. We therefore elected to analyze each RCT individually and to then discuss their common **and their differing findings** on the background of the pertaining limitations.

The first study [29] compared posterior endoscopic foraminotomy to microsurgical anterior decompression

and fusion (ACDF; using a stand-alone poly-ether-etherketone [PEEK]-cage) for the treatment of lateral cervical disc herniations. This study was randomized and included 200 patients with a follow-up of 2 years and a follow-up rate of 88 %. The key findings were a reduced operating time (28 vs. 68 minutes on average) and a faster return to work (19 vs. 34 days) with the endoscopic technique. **Clinical outcome and complication rates** were not significantly different between the techniques.

The second study [30] compared endoscopic interlaminar and transforaminal lumbar discectomy to the conventional microsurgical technique (microdiscectomy) in lumbar disc herniations, irrespective of their location (median, lateral, or extraforaminal). This study was randomized and included 200 patients with a follow-up of 2 years and a follow-up rate of 89 %. The key findings were a reduced operating time (22 vs. 43 minutes on average) and a faster return to work (25 vs. 49 days) with the endoscopic technique. Clinical outcome was not significantly different between the techniques, but there were more progressions to fusion and more light complications with the traditional technique.

The third study [31] compared the same techniques as in the second study, but for recurrent lumbar disc herniations after a previous conventional **microdiscectomy**. This study was randomized and included 100 patients with a follow-up of 2 years and a follow-up

Table 2

Study Outcome Criteria and Results

Authors/year/ herniation type	Procedure	Operating time (min)/ blood loss (ml)	Postoperative pain/pain medication	Clinical outcome criteria (preop to last FU)	Radiological outcome criteria	Max. hospital stay/return to work (days)	Reoperations	Complications	Poor clinical outcome
Ruetten et al., 2008 cervical lateral	endoscopic posterior foraminotomy	28 (19–50)/ 0	«significantly less» (no detailed data given)	VAS neck: 17 to 16 VAS arm: 84 to 7 NASSpain: 4.1 to 1.4 NASS neurology: 3.2 to 1.6 Hilibrand: 0 to 86 (good/ excellent)	advancing degeneration of index disc: 21 (24%) kyphosis progression or instability: 0	dna/19	6 overall (6.6%) 3 for persistent arm pain 3 for recurrent herniation	transient dermatomal hypoesthesia: 3 serious complications: 0	9
	microsurgical ACDF (PEEK cage, no plate)	68 (48–105)/ <10		VAS neck: 15 to 17 VAS arm: 81 to 8 NASSpain: 4.3 to 1.5 NASS neurology: 3.4 to 1.6 Hilibrand: 0 to 81 (good/excellent)	incomplete fusion: 17 (18%) subsidence (3 mm max): 5 (5.8%)	dna/34	4 overall (4.8%) 3 for persistent arm pain 1 for implant failure	transient difficulty swallowing: 3 surface hematoma: 1 distorted scar: 1 serious complications: 0	8
Ruetten et al., 2008 lumbar (median, lateral, extra-foraminal)	endoscopic interlaminar or transforaminal sequestrectomy	22 (13–46)/ 0	«significantly less» (no detailed data given)	VAS back: 19 to 11 VAS leg: 75 to 8 ODI: 75 to 20 NASS pain: 4.6 to 2.1 NASS neurology: 3.1 to 2.1	dna	dna/25	9 overall (9.9%) 6 for recurrent herniation 2 for repeated recurrence 1 fusion for progressive LBP	transient postoperative dysesthesia: 3 serious complications: 0	13 overall (not differentiated between groups)
	microsurgical sequestrectomy (paramedian or lateral)	43 (34–72)/ 45 (5–235)		VASback: 15 to 18 VAS leg: 71 to 9 ODI: 73 to 24 NASSpain: 4.2 to 2.3 NASS neurology: 2.9 to 1.9	dna	dna/49	10 overall (11.5%) 5 for recurrent herniation 5 fusions for progressive LBP	transient postoperative dysesthesia: 5 postoperative bleeding: 2 delayed wound- healing: 2 soft tissue infection: 1 transient urinary retention: 3 serious complications: 0	13 overall (not differentiated between groups)
Ruetten et al., 2008 lumbar recurrent	endoscopic interlaminar or transforaminal sequestrectomy	24 (14–43)/ 0	«significantly less» (no detailed data given)	VAS back: 14 to 15 VASleg: 79 to 8 ODI: 80 to 20 NASSpain: 4.3 to 2.1 NASS neurology: 2.5 to 2.1	dna	dna/28	5 overall (11.1%) 3 for recurrent herniation 2 for persistent leg pain	dural tears: 1 transient postoperative dysesthesia: 2 wound complications: 0 serious complications: 6%	9 overall (not differentiated between groups)
	microsurgical sequestrectomy (paramedian approach)	58 (39–91)/ 41 (10–205)		VASback: 15 to 14 VAS leg: 85 to 10 ODI: 84 to 21 NASS pain: 4.5 to 2.1 NASS neurology: 2.3 to 2.3	dna	dna/52	5 overall (11.9%) 2 for recurrent herniation 1 for persistent leg pain 2 fusions for progressive LBP	dural tears: 3 transient postoperative dysesthesia: 5 delayed wound healing: 2 soft tissue infection: 1 serious complications: 21%	9 overall (not differentiated between groups)

Table 2 (cont.)

Study Outcome Criteria and Results

Authors/year/ herniation type	Procedure	Operating time (min)/ blood loss (ml)	Postoperative pain/pain medication	Clinical outcome criteria (preop to last FU)	Radiological outcome criteria	Max. hospital stay/return to work (days)	Reoperations	Complications	Poor clinical outcome
Ruetten et al., 2008 cervical	anterior transfascial endoscopic decompression	32 (18-51)/0	dna	VAS neck: 18 to 15 VAS arm: 82 to 8 NASS pain: 4.2 to 1.5 NASS neurology: 3.1 to 1.8 Hilibrand: 0 to 48 (good/ excellent)	progression of adjacent disc degeneration: 3 kyphosis increase index segment: 6 (5°) index disc space height: 5.3 to 4.1 mm no spontaneous fusion progressive degeneration index disc: 21 (24%)	3/at 3 months 84%	4 overall (7.4%) 2 for recurrent herniation 2 for persistent arm pain	transient difficulty swallowing: 2 superficial hematoma: 0 serious complications: 0	11 overall (not differentiated between groups)
	microsurgical ACDF (PEEK cage, no plate)	62 (41-102)/ <10	dna	VAS neck: 13 to 14 VAS arm: 79 to 10 NASS pain: 4.4 to 1.6 NASS neurology: 3.2 to 1.6 Hilibrand: 0 to 43 (good/excellent)	progression of adjacent disc degeneration: 9 kyphosis increase index segment: 4 (5.9°) index disc space height: 6.1 to 5.0 mm incomplete fusion: 9 (18.8%)	7/at 3 months 63% (sig)	3 overall (6.1%) 2 for persistent arm pain 1 repeat ACDF for subsidence	transient difficulty swallowing: 5 superficial hematoma: 2 serious complications: 0	11 overall (not differentiated between groups)
Lee et al., 2010 lumbar recurrent	endoscopic transforaminal sequestrectomy and disc decompression	46 ± 11/ dna	dna	VAS back: 7.0 to 2.9 VAS leg: 8.4 to 2.9 ODI: 62 to 21	secondary instability: 0 index disc space height: 34 to 32% sagittal rotational angle: 9 to 9.5° multIHtus muscle volume: 713mm2to717 mmr2	0.9 ± 0.5	2 overall (8%) 1 for persistent leg pain 1 fusion for repeated recurrence	persistent leg pain: 1	dna
	microsurgical sequestrectomy (paramedian approach)	74 ± 26	dna	VAS back: 5.4 to 3.1 VAS leg: 8.6 to 3.5 ODI: 63 to 18	secondary instability: 1 index disc space height: 35 to 32% sagittal rotational angle: 8.2° to 9.9° multifidus muscle volume: 730 mm2 to 705 mm2	3.8 ± 1.4	1 overall (3.4%) 1 fusion for repeated recurrence	dural tears: 2 incomplete cauda equine syndrome: 1	dna

Note. Ref Nr. refers to the specific reference number in the manuscript bibliography, FU = follow up, ACDF = anterior cervical decompression and fusion, PF.FK = poly-ether-ether-ketone, dna = data not available, VAS = visual analog scale, NASS = North American Spine Society, ODI = Oswestry Disability Index. For each trial, data on the endoscopic procedure are displayed in the first line, data on the comparison procedure in the second line.

rate of 87 %. **The key findings were a reduced operating time (24 vs. 58 minutes on average) and a faster return to work (28 vs. 52 days) with the endoscopic technique.** Clinical outcome was **not significantly different** between the techniques, but there were more serious complications with the traditional technique.

The fourth study [32] compared endoscopic anterior (transdiscal) decompression to microsurgical ACDF using a stand-alone PEEK-cage in cervical disc herniations.

This study was randomized and included 120 patients with a follow-up of 2 years and a follow-up rate of 86 %. The key findings were a reduced operating time (32 vs. 36 minutes on average) and a higher percentage of patients having returned to work after 3 months (84 vs. 63 %) with the endoscopic technique. Clinical outcome and complication rates were not significantly different between the techniques.

The fifth study [37] compared transforaminal endoscopic lumbar discectomy to lumbar microdiscectomy in recurrent disc herniations. This study had a retrospective, non-randomized design and included 54 patients with an average follow-up of 34 months. The key findings were a reduced operating time (46 vs. 74 minutes on average) and a shorter average stay in hospital (0.9 vs. 3.8 days) with the endoscopic technique. Clinical outcome and complication rates were not significantly different between the techniques.

What are the common findings among these 5 studies?

All 5 studies demonstrated significant improvement in their clinical target criteria between preoperative and the different time points until final follow-up. No study showed significant differences in these target criteria between the endoscopic technique and the respective standard technique. All 5 studies had significantly shorter operating times for the endoscopic technique compared to the respective standard technique. The reoperation rates were comparable between groups with a possible tendency towards slightly higher reoperation rates with the endoscopic technique in 2 studies [29, 37]. The radiological target criteria did not show any clinically relevant differences between the endoscopic and the standard technique groups with the exception of one segmental instability in the standard technique group that led to a fusion surgery in the study by Lee et al. [37].

What are the differences between the endoscopic and the standard techniques with regards to complications and reoperations?

In all 5 studies, there were fewer complications reported with the endoscopic techniques compared to the standard techniques. In 2 of the 5 studies, these differences reached statistical significance [30, 31].

Of these 2 studies, the one on primary lumbar disc herniations showed a higher rate of revision fusion procedures for progressive low back pain (5 vs. one) [30], whereas the study on recurrent lumbar disc herniations showed a difference in serious complications of 21 % vs. 6 %, both in favor of the endoscopic approach [31]. The study comparing anterior cervical discectomy and fusion (ACDF) to endoscopic anterior transdiscal decompression for cervical disc herniations found less postoperative difficulty with swallowing (5 vs. 2) in the endoscopic group without reaching statistical significance [32].

What appear to be the benefits of the endoscopic technique in these 5 studies?

Obvious benefits are shorter operating times and less blood loss (even if not statistically evaluated) in all 5 studies. Three of the 5 studies claim significantly less pain at the surgical site immediately postoperative and less use of pain medication [29–31], but detailed data are not contained in the respective publications. These observations are paralleled by a shorter hospital stay in one [37] and a faster return to work in the 4 other studies [29–32]. As described above, there were fewer complications with the endoscopic technique as compared to the standard techniques in all of the 5 studies and lower rates of revision fusion surgeries in one study.

What were the advantages of the standard technique in these 5 studies?

The standard techniques appear to have an advantage with regards to the rates of recurrent herniations and repeated recurrent herniations in cervical as well as in lumbar disc surgery, even though statistical significance was not shown in these studies [29, 31, 32].

What do the radiological target parameters tell us?

In the 2 trials on cervical disc herniations, radiologically uncertain fusions were observed in almost a fifth of the ACDF cases at 24 months without translating to reduced clinical success, which is not a surprising finding [29, 32]. A much more interesting observation is that no increased segmental kyphosis was observed after the anterior endoscopic transdiscal approach in the second study when compared to ACDF and that there appeared to be less progression in pre-existing adjacent level disc degeneration with the endoscopic technique [32].

When comparing posterior endoscopic foraminotomy to ACDF, Ruetten et al. [29] found a progression in the radiological degeneration of the index disc in 24 % of cases without concomitant progression of segmental kyphosis or creation of new segmental instability. The study by Lee et al. [37] showed a pronounced (and statistically significant) decrease in the index disc space

height as well as a stronger increase in the sagittal rotational angle when comparing microdiscectomy to a transforaminal endoscopic technique.

Discussion

The limitations of this review should be examined first. The 5 studies selected by means of our search strategy and exclusion process, while all employing the same full-endoscopic technique, are heterogeneous with regards to spinal region, type of approach and **herniation**, as well as comparison procedure. It must also be taken into account that the surgeons who performed the RCTs and the CSs evaluated in our review are highly experienced and specialized in the standard techniques as well as in the endoscopic techniques that they studied. The data that found entry into these studies certainly do not originate from the first few hundred cases that these investigators performed. The results obtained in these trials can therefore not be directly translated to what other spinal therapists at an earlier stage of their individual learning curves can expect to achieve.

A second, yet very important, limitation is the fact that all 4 RCTs in this review were performed by the same group of investigators and at the same institution. It will remain to be seen whether other endoscopic spine surgeons in different settings and with a **different training** background will be able to duplicate these results.

In the context of these limitations, however, our review finds benefits to the patient with these modern, full-endoscopic techniques. Most importantly and at least in the 5 studies that could be considered for this review, these benefits do not appear to come at the cost of increased complication rates or lesser efficacy. Shorter operating times and less postoperative surgical site pain translate to a shorter hospital stay and may lead to a faster return to work.

While the claims of less postsurgical pain with the endoscopic technique made by 3 of the 5 trials [29–31] appear credible based on the access trauma of the comparison procedures, it is a severe shortcoming of these 3 studies that no clear data are contained in the published manuscripts. There was a statistically not significant higher rate of reoperations for recurrence in some of the studies, but a claim that these rates would have become significant with larger numbers is difficult to make at around 100 patients per group. They may however become significant with lesser surgeon experience, which should be considered when extrapolating from these studies to the personal case series and the same is most probably true for complication rates. A solid experience with the standard techniques should therefore remain the basis on which these endoscopic techniques can be mastered step by step.

While this is not uncommon in clinical medicine, it is **nevertheless regrettable that with so many publications on endoscopic spine surgery, so few controlled studies are available that compare an established standard procedure to a modern full-endoscopic procedure.** Beyond the mere paucity of suitable studies, the fact that most of the available controlled studies originate from one single group of very specialized researchers is a limitation on the generalizability of the clinical results obtained. Having said that and considering the poor quality of data that are the foundation for the establishment of microdiscectomy as the de facto gold standard over open discectomy, the results from the trials discussed here are already a big step in the right direction.

With most of the relevant studies in this field having been published within the past 4 years, there are only few previous reviews on this topic. The most notable one is the paper by Nellensteijn et al. [12] submitted in 2009 and published in 2010. This review, however, is very different from ours in a number of respects, some of which have already been mentioned in the introduction. **It does not consider cervical disc herniations or other endoscopic approaches other than the lumbar transforaminal approach.** The literature search for this review also ends with 2008, so that 3 of the 5 papers that we were able to consider did not find entry into the authors' evaluation. The other review by the same group and also published in 2010 focuses exclusively on transforaminal endoscopic surgery for spinal stenosis, which certainly is not an indication that is frequently treated by means of transforaminal endoscopic techniques [12].

Conclusion

In summary, there is good quality evidence, but from predominantly only one group, that experienced surgeons can achieve the same clinical results in symptomatic cervical and lumbar disc herniations with full endoscopic procedures as with standard microsurgical procedures.

In experienced hands, these results do not appear to come at the cost of a higher complication rate or more severe complications.

References

1. Saberski L. R. A retrospective analysis of spinal canal endoscopy and laminectomy outcomes data / L. R. Saberski // Pain Physician. — 2000. — Vol. 3. — P. 193–196.
2. Kambin P. Arthroscopic microdiscectomy / P. Kambin // Arthroscopy. — 1992. — Vol. 8. — P. 287–295.
3. Kovac D. Automated endoscopic percutaneous discectomy in the treatment of lumbar disk hernia / D. Kovac // Lijec Vjesn. — 1991. — Vol. 113. — P. 158–161.
4. Leu H. J. 10 years of percutaneous disk surgery: results and

- developments / H. J. Leu, A. Schreiber // *Schweiz Rundsch Med Prax.* — 1989. — Vol. 78. — P. 1434–1439.
5. Suezawa Y. Percutaneous nucleotomy with discoscopy. 7 years' experience and results / Y. Suezawa, A. Schreiber // *Z. Orthop. Ihre Grenzgeb.* — 1988. — Vol. 126. — P. 1–7.
 6. Schreiber A. Transdiscoscopic percutaneous nucleotomy in disk herniation / A. Schreiber, Y. Suezawa // *Orthop. Rev.* — 1986. — Vol. 15. — P. 35–38.
 7. Hausmann B. Nucleoscope. Instrumentarium for endoscopy of the intervertebral disc space / B. Hausmann, R. Forst // *Arch. Orthop. Trauma Surg.* — 1983. — Vol. 102. — P. 57–59.
 8. Gibson J. Surgical interventions for lumbar disc prolapse / J. Gibson, G. Waddell // *Spine.* — 2007. — Vol. 32. — P. 1735–1747.
 9. Effectiveness of microdiscectomy for lumbar disc herniation: A randomized controlled trial with 2 years of follow-up / H. Osterman, S. Seitsalo, J. Karppinen, A. Malmivaara // *Spine.* — 2006. — Vol. 31. — P. 2409–2414.
 10. Comparison of surgical outcomes between macro discectomy and micro discectomy for lumbar disc herniation: A prospective randomized study with surgery performed by the same spine surgeon / Y. Katayama, Y. Matsuyama, H. Yoshihara [et al.] // *Journal Spinal Disorders Techniques.* — 2006. — Vol. 19. — P. 344–347.
 11. A controlled study of microsurgical versus standard lumbar discectomy / L. Henriksen, K. Schmidt, V. Eskesen, E. Jantzen // *Br. J. Neurosurg.* — 1996. — Vol. 10. — P. 289–293.
 12. Transforaminal endoscopic surgery for symptomatic lumbar disc herniations: a systematic review of the literature / J. Nellensteijn, R. Ostelo, R. Bartels [et al.] // *Eur. Spine J.* — 2010. — Vol. 19. — P. 181–204.
 13. Less systemic cytokine response in patients following microendoscopic versus open lumbar discectomy / T. J. Huang, R. W. Hsu, Y. Y. Li, C. C. Cheng // *J. Orthop. Res.* — 2005. — Vol. 23. — P. 406–411.
 14. Microendoscopic lumbar discectomy versus open surgery: an intraoperative EMG study / U. Schick, J. Dohnert, A. Richter [et al.] // *Eur. Spine J.* — 2002. — Vol. 11. — P. 20–26.
 15. Tubular discectomy vs conventional microdiscectomy for sciatica: a randomized controlled trial / M. P. Arts, R. Brand, M. E. van den Akker [et al.] // *JAMA.* — 2009. — Vol. 302. — P. 149–158.
 16. Higher risk of dural tears and recurrent herniation with lumbar micro-endoscopic discectomy / M. Teli, A. Lovi, M. Brayda-Bruno [et al.] // *Eur. Spine J.* — 2010. — Vol. 19. — P. 443–450.
 17. A critical review of the American Pain Society Clinical practice guidelines for interventional techniques: Part 2. Therapeutic interventions / L. Manchikanti, S. Datta, S. Gupta [et al.] // *Pain Physician.* — 2010. — Vol. 13. — P. E215–E264.
 18. An algorithmic approach for clinical management of chronic spinal pain / L. Manchikanti, S. Helm, V. Singh [et al.] // *Pain Physician.* — 2009. — Vol. 12. — P. E225–E264.
 19. Reassessment of evidence synthesis of occupational medicine practice guidelines for interventional pain management / L. Manchikanti, V. Singh, R. Derby [et al.] // *Pain Physician.* — 2008. — Vol. 11. — P. 393–482.
 20. Systematic review of effectiveness and complications of adhesiolysis in the management of chronic spinal pain: an update / A. M. Trescot, P. Chopra, S. Abdi [et al.] // *Pain Physician.* — 2007. — Vol. 10. — P. 129–146.
 21. Interventional techniques: Evidencebased practice guidelines in the management of chronic spinal pain / M. V. Boswell, A. M. Trescot, S. Datta [et al.] // *Pain Physician.* — 2007. — Vol. 10. — P. 7–111.
 22. Role of adhesiolysis in the management of chronic spinal pain: A systematic review of effectiveness and complications / P. Chopra, H. S. Smith, T. R. Deer, R. C. Bowman // *Pain Physician.* — 2005. — Vol. 8. — P. 87–100.
 23. Interventional techniques in the management of chronic spinal pain: Evidencebased practice guidelines / M. V. Boswell, R. V. Shah, C. R. Everett [et al.] // *Pain Physician.* — 2005. — Vol. 8. — P. 1–47.
 24. Minisurgical approach for spinal endoscopy in the presence of stenosis of the sacral hiatus / S. Helm 2nd, J. D. Gross, K. G. Varley [et al.] // *Pain Physician.* — 2004. — Vol. 7. — P. 323–325.
 25. Spinal endoscopic adhesiolysis in the management of chronic low back pain: a preliminary report of a randomized, double-blind trial / L. Manchikanti, J. J. Rivera, V. Pampati [et al.] // *Pain Physician.* — 2003. — Vol. 6. — P. 259–267.
 26. Transforaminal posterolateral endoscopic discectomy with or without the combination of a low-dose chymopapain: a prospective randomized study in 280 consecutive cases / T. Hoogland, M. Schubert, B. Miklitz, A. Ramirez // *Spine.* — 2006. — Vol. 31. — P. E890–E897.
 27. Interleukin-6 protein expression is more important than interleukin-6 mRNA levels in assessing surgical invasiveness / T. J. Huang, Y. Y. Li, Y. J. Weng [et al.] // *J. Surgical Research.* — 2007. — Vol. 142. — P. 53–58.
 28. Righesso O. Comparison of open discectomy with microendoscopic discectomy in lumbar disc herniations: results of a randomized controlled trial / O. Righesso, A. Falavigna, O. Avanzi // *Neurosurgery.* — 2007. — Vol. 61. — P. 545–549.
 29. Full-endoscopic cervical posterior foraminotomy for the operation of lateral disc herniations using 5.9-mm endoscopes: a prospective, randomized, controlled study / S. Ruetten, M. Komp, H. Merk, G. Godolias // *Spine.* — 2008. — Vol. 33. — P. 940–948.
 30. Full-endoscopic interlaminar and transforaminal lumbar discectomy versus conventional microsurgical technique: a prospective, randomized, controlled study / S. Ruetten, M. Komp, H. Merk, G. Godolias // *Spine.* — 2008. — Vol. 33. — P. 931–939.
 31. Recurrent lumbar disc herniation after conventional discectomy: a prospective, randomized study comparing full-endoscopic interlaminar and transforaminal versus microsurgical revision / S. Ruetten, M. Komp, H. Merk, G. Godolias // *J. Spinal Disord. Tech.* — 2009. — Vol. 22. — P. 122–129.
 32. Full-endoscopic anterior decompression versus conventional anterior decompression and fusion in cervical disc herniations / S. Ruetten, M. Komp, H. Merk, G. Godolias // *Int. Orthop.* — 2009. — Vol. 33. — P. 1677–1682.
 33. Analysis of the results of endoscopic surgical treatment for hernias of the lumbosacral disks / S. K. Akshulakov, T. T. Kerimbaev, G. T. Sekenova, G. S. Makimova // *Zh Vopr Neurokhir im N. N. Burdenko.* — 2005. — № 4. — P. 18–21.
 34. Microendoscopic discectomy, a less traumatic procedure for lumbar disk herniation / Z. Chao, Z. Yue, C. Tong-wei [et al.] // *Chin J. Traumatol.* — 2007. — Vol. 10. — P. 311–314.
 35. Kang Y. Sagittal spinal alignment in patients with lumbar disc herniation after endoscopic discectomy / Y. Kang, K. Endo // *Journal of Tokyo Medical University.* — 2009. — Vol. 67. — P. 431–438.
 36. Lee D. Y. Learning curve for percutaneous endoscopic lumbar discectomy / D. Y. Lee, S. H. Lee // *Neurol. Med. Chir. (Tokyo).* — 2008. — Vol. 48. — P. 383–388.
 37. Comparison of percutaneous endoscopic lumbar discectomy and open lumbar microdiscectomy for recurrent disc herniation / D. Y. Lee, C. S. Shim, Y. Ahn [et al.] // *J. Korean Neurosurg. Soc.* — 2009. — Vol. 46. — P. 515–521.
 38. Long-term outcomes of percutaneous lumbar discectomy and microendoscopic discectomy for lumbar disc herniation / W. G. Liu, X. T. Wu, J. Min [et al.] // *Zhonghua Yi Xue Za Zhi.* — 2009. — Vol. 89. — P. 750–753.
 39. Mayer H. M. Percutaneous endoscopic discectomy: surgical

- technique and preliminary results compared to microsurgical discectomy / H. M. Mayer, M. Brock // *J. Neurosurg.* — 1993. — Vol. 78. — P. 216–225.
40. Objective assessment of reduced invasiveness in MED: Compared with conventional one-level laminotomy / R. Sasaoka, H. Nakamura, S. Konishi [et al.] // *European Spine Journal.* — 2006. — Vol. 15. — P. 577–582.
41. The efficacy of microendoscopic discectomy in reducing iatrogenic muscle injury / D. A. Shin, K. N. Kim, H. C. Shin, Yoon do H. // *J. Neurosurg. Spine.* — 2008. — Vol. 8. — P. 39–43.
42. Low-back pain following surgery for lumbar disc herniation: A prospective study / T. Toyone, T. Tanaka, D. Kato, R. Kaneyama // *J. Bone Joint Surgery.* — 2004. — Vol. 86-A. — P. 893–896.
43. Xiao L. Z. A comparative study about curative effect of two different therapies to the lumbar disc herniation [Chinese] / L. Z. Xiao, D. R. Zhang, J. Jiang // *Chinese Journal of Clinical Rehabilitation.* — 2002. — Vol. 6. — P. 552–553.
44. Identifying pathogens of spondylodiscitis: percutaneous endoscopy or CTguided biopsy / S. C. Yang, T. S. Fu, L. H. Chen [et al.] // *Clin. Orthop. Related Research.* — 2008. — Vol. 466. — P. 3086–3092.
45. Microendoscopic discectomy, a less traumatic procedure for lumbar disk herniation / C. Zhang, Y. Zhou, T. W. Chu [et al.] // *Chin. J. Traumatol. (Engl. Ed.).* — 2007. — Vol. 10. — P. 311–314.
46. Traumatic responses following microendoscopic discectomy: clinical analysis of 44 patients / C. Zhang, T. W. Chu, Y. Zhou [et al.] // *Zhonghua Yi Xue Za Zhi.* — 2006. — Vol. 86. — P. 3039–3042.
47. Minimally invasive strategies and options for farlateral lumbar disc herniation / Y. Zhou, C. Zhang, J. Wang [et al.] // *Chin. J. Traumatol.* — 2008. — Vol. 11. — P. 259–266.

Article submitted on 13.05.2014

СОВРЕМЕННОЕ СОСТОЯНИЕ ЭНДОСКОПИЧЕСКОЙ ХИРУРГИИ МЕЖПОЗВОНКОВОГО ДИСКА: ОБЗОР КОНТРОЛИРУЕМЫХ ИССЛЕДОВАНИЙ СРАВНЕНИЯ ЭНДОСКОПИЧЕСКИХ И СТАНДАРТНЫХ ОПЕРАЦИЙ ДЛЯ ЛЕЧЕНИЯ ГРЫЖ МЕЖПОЗВОНКОВЫХ ДИСКОВ

Кристоф Биркенмайер¹, Мартин Комп², Хансйорг Ф. Лой³, Бернд Вегенер¹, Себастиан Рюттен²

¹ Ортопедическое отделение, Мюнхенский Университет (LMU), Кампус Гроссхадерн, Мюнхен. Германия

² Хернесская больница Святой Анны, Университет Виттен/Хердеке, Херне. Германия

³ Центр позвоночника Бетания, Больница Бетания, Цюрих. Швейцария